

NEXT ON YOUR  
LITTLE PINK JOURNEY...

addyi<sup>®</sup>  
(flibanserin)  
100mg tablets

phlRX WILL TEXT YOU &  
CONFIRM YOUR PAYMENT

CLICK THE LINK TO PAY (AND GET LOWEST COPAY)



TRACK YOUR SHIPMENT

YOU'LL RECEIVE TRACKING ONCE SHIPPED

QUESTIONS? CONTACT PATIENT SUPPORT: 855-902-2779 X 3



TAKE YOUR ADDYI DAILY,  
ONLY AT BEDTIME

It is important to take Addyi as directed by your healthcare provider to help decrease the risks of low blood pressure, fainting, accidental injury and sleepiness.



WATCH FOR SIGNS

AFTER A FEW WEEKS\* ON ADDYI, YOU MAY NOTICE:

- SEXUAL FANTASIES
- WANTING TO HAVE SEX AGAIN
- RENEWED INTEREST IN A PARTNER'S INITIATION

\*Some women may see results in as early as 4 weeks.

Addyi should be discontinued if there is no improvement after 8 weeks.



#### INDICATION

##### What is ADDYI (add-ee) (flibanserin) Tablets?

ADDYI is a prescription medicine used to treat hypoactive (low) sexual desire disorder (HSDD) in women who have not gone through menopause, who have not had problems with low sexual desire in the past, and who have low sexual desire no matter the type of sexual activity, the situation, or the sexual partner. Women with HSDD have low sexual desire that is troubling to them. Their low sexual desire is not due to: a medical or mental health problem, problems in the relationship, medicine or other drug use. ADDYI is not for use for the treatment of HSDD in women who have gone through menopause, in men or in children. ADDYI is not for use to enhance sexual performance.

#### Selected Important Safety Information

Your risk of severe low blood pressure and fainting is increased if you drink alcohol close in time to your ADDYI dose, take certain prescription, OTC or herbal medications, or have liver problems.

See Important Safety Information, including Boxed Warning regarding severe low blood pressure and fainting in certain settings, on back and Full Prescribing Information and Medication Guide at [addyi.com/pi](http://addyi.com/pi).



addyi<sup>®</sup>  
(flibanserin)  
100mg tablets

#### IMPORTANT SAFETY INFORMATION

What is the most important information I should know about ADDYI? Your risk of severe low blood pressure and fainting (loss of consciousness) is increased if you take ADDYI and:

- drink alcohol close in time to when you take your ADDYI dose.
- Wait at least 2 hours after drinking 1 or 2 standard alcoholic drinks before taking ADDYI at bedtime.

Examples of 1 standard alcoholic drink include: one 12-ounce regular beer, 5 ounces of wine, 1.5 ounces of distilled spirits or shot

- If you drink 3 or more standard alcoholic drinks in the evening, skip your ADDYI dose at bedtime.

- After you have taken your ADDYI at bedtime, do not drink alcohol until the following day.

- take certain prescription medicines, over-the-counter medicines, or herbal supplements. Do not take or start taking any prescription medicines, over-the-counter medicines, or herbal supplements while taking ADDYI until you have talked with your doctor. Your doctor will tell you if it is safe to take other medicines or herbal supplements while you are taking ADDYI.

- have liver problems. Do not take ADDYI if you have liver problems.

If you take ADDYI and you feel lightheaded or dizzy, lie down right away. Get emergency medical help or ask someone to get emergency medical help for you if the symptoms do not go away or if you feel like you could faint (lose consciousness). If you faint (lose consciousness), tell your doctor as soon as you can.

#### Who should not take ADDYI?

##### Do not take ADDYI if you:

- take certain medicines. Taking ADDYI with certain other medicines can increase the amount of ADDYI in your blood and cause severe low blood pressure, fainting (loss of consciousness), and sleepiness.
- Do not take ADDYI if you are taking any of the following medicines:
  - Certain medicines used to treat HIV-1 infection
  - Certain medicines that you take by mouth used to treat fungal infections
  - Certain antibiotics
  - Certain medicines used to treat Hepatitis C infection
  - Certain medicines used to treat high blood pressure, chest pain (angina), or other heart problems
  - Nefazodone: a medicine used to treat depression

Ask your doctor or pharmacist if you are not sure if you take any of the medicines listed above. These are examples of the medicines that you should not take if you are taking ADDYI. Tell your doctor about all the medicines you take before you start taking ADDYI

- have liver problems

are allergic to flibanserin or any of the ingredients in ADDYI. See Medication Guide for the full list of ingredients.

#### What should I tell my doctor before taking ADDYI?

Before you take ADDYI, tell your doctor about all of your medical conditions, including if you:

- drink alcohol, use drugs, or have a history of alcohol or drug abuse
- have ever had depression or other mental health problems
- have low blood pressure or a medical condition that can cause low blood pressure

- are pregnant or plan to become pregnant. It is not known if ADDYI will harm your unborn baby.

- are breastfeeding or plan to breastfeed. It is not known if ADDYI passes into your breast milk. You and your doctor should decide if you will take ADDYI or breastfeed. You should not do both.

Tell your doctor if you have had an allergic reaction such as hives, itching, or trouble breathing during or after receiving a dose of ADDYI.

Tell your doctor about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. ADDYI can affect the way other medicines work, and other medicines can affect the way ADDYI works, and can cause serious side effects.

#### What should I avoid while taking ADDYI?

- Do not drink alcohol close to the time you take your ADDYI dose because this increases your risk of severe low blood pressure and fainting (loss of consciousness).

- Do not drive, operate machinery, or do things that require clear thinking until at least 6 hours after you take ADDYI and until you know how ADDYI affects you.

- Do not drink grapefruit juice if you take ADDYI. Drinking grapefruit juice during your treatment with ADDYI increases your risk of severe low blood pressure and fainting (loss of consciousness).

- You should not take the herbal supplements St. John's Wort, ginkgo, or resveratrol or certain over-the-counter medicines such as cimetidine until you talk to your doctor. Taking ADDYI with these herbal supplements and over-the-counter medicines may increase your risk of low blood pressure, fainting (loss of consciousness), and sleepiness.

#### What are the possible side effects of ADDYI?

ADDYI can cause serious side effects, including:

- Sleepiness is a common side effect of ADDYI and can be serious. Taking ADDYI can increase your risk of sleepiness if taken during waking hours, if you drink alcohol, or take certain medicines or herbal supplements.
- Low blood pressure and fainting (loss of consciousness) can happen when you take ADDYI even if you do not drink alcohol or take other medicines or herbal supplements. Your risk of low blood pressure and fainting (loss of consciousness) is increased if ADDYI is taken during waking hours, if you drink alcohol within 2 hours of taking ADDYI, or if you take certain medicines or herbal supplements.

#### The most common side effects of ADDYI include:

- Dizziness
- Difficulty falling asleep or staying asleep
- Nausea
- Dry mouth
- Tiredness

These are not all of the possible side effects of ADDYI. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

See Indication on reverse side and accompanying Full Prescribing Information and Medication Guide, including Boxed Warning regarding severe low blood pressure and fainting in certain settings at [addyi.com/pi](http://addyi.com/pi).

This information does not take the place of talking with your doctor.